



Review Article

Simulation in paediatric urology and surgery, part 2: An overview of simulation modalities and their applications

R.M. Nataraja ^{a,b}, N. Webb ^c, P.J. Lopez ^{c,d,e}

^aDepartment of Paediatric Surgery, Monash Children's Hospital, Melbourne Australia

^bDepartment of Paediatrics, School of Clinical Sciences, Faculty of Medicine, Nursing and Health Sciences, Monash University, Melbourne Australia

^cDepartment of Paediatric Urology, Monash Children's Hospital, Melbourne Australia

^dDepartment of Urology, Hospital Exequiel Gonzalez Cortes, Santiago, Chile

^eClinica Alemana, Santiago, Chile

Correspondence to: R.M. Nataraja, Monash Children's Hospital, Department of Paediatric Surgery, 246 Clayton Road, Clayton, Melbourne 3168, Australia

ram.nataraja@monashhealth.org (R.M. Nataraja)

Keywords

Surgical simulation; Simulation based medical education (SBME); Urology simulation; Paediatric surgery simulation; Paediatric urology simulation

Received 31 August 2017
Accepted 29 December 2017
Available online xxx

Summary

Surgical training has changed radically in the last few decades. The traditional Halstedian model of time-bound apprenticeship has been replaced with competency-based training. In our previous article, we presented an overview of learning theory relevant to clinical teaching; a summary for the busy paediatric surgeon and urologist. We introduced the concepts underpinning current changes in surgical education and training. In this next article, we give an overview of the various modalities of surgical

simulation, the educational principles that underlie them, and potential applications in clinical practice. These modalities include; open surgical models and trainers, laparoscopic bench trainers, virtual reality trainers, simulated patients and role-play, hybrid simulation, scenario-based simulation, distributed simulation, virtual reality, and online simulation. Specific examples of technology that may be used for these modalities are included but this is not a comprehensive review of all available products.

Introduction

Healthcare education has evolved rapidly in the past two decades, with changes especially evident in surgical training. There has been a shift from the traditional Halstedian model of time-bound apprenticeship to one of competency-based training [1]. Simulation methodology is now an integral part of training in various surgical subspecialties, especially for technical skills learning. However, it is important to recognize that the role of simulation in surgical education is broader than merely technical skill acquisition. Many adverse incidents in surgical practice arise from or involve failures in domains such as communication, teamwork, and situational awareness, rather than technical expertise [2–4]. Simulation can be employed to promote the learning, practice, refinement, and assessment of both technical and non-technical skills in a patient-safe environment. In addition to allowing learners to make mistakes without adverse patient impacts, simulation also allows for specific rehearsal of rare or unique situations.

In part 1 of this simulation-based medical education (SBME) overview, we introduced the various educational theory concepts that underpin learning clinical environment. The aim

of this second part is to provide an overview of the various simulation modalities that can be used to apply these principles and theories.

Open surgery models and trainers

The use of simulators in surgical training is not a novel concept, having been used in many forms for centuries [5]. The majority of these are relatively simple 'part-task' trainers, familiar to most surgeons. Despite their reductionist simplicity, they promote the acquisition of basic skills that form the foundations of surgical expertise. These types of simulators rely on the principles of deliberate practice and feedback to foster development of competency. Part task trainer (PTT) simulators can include cadaveric, synthetic, or animal models that simulate part of a surgical intervention [6]. The use of cadaveric tissue is becoming less common as the availability of alternate bench training models with educational equivalence increases [7]. In some environments, access to and use of biologic tissues is severely limited.

Basic surgical skills (BSS) courses using PTTs are an integral element of early surgical training in many countries, possible even in resource-constrained environments [8]. These PTTs include hooks positioned on a wooden

<https://doi.org/10.1016/j.jpuro.2017.12.009>

1477-5131/© 2018 Journal of Pediatric Urology Company. Published by Elsevier Ltd. All rights reserved.

board for hand tying sutures, synthetic skin with a laceration for suturing, bowel, or vascular anastomosis using porcine aorta. These courses are often the first introduction of surgical trainee to bench training simulation.

With the introduction of more readily available additive manufacturing techniques or 3D printing, there has been a rapid increase in the number of models for surgical training [9,10]. Hybrid models of surgically altered bovine or porcine tissue, combined with 3D printed chest or abdomen covered in synthetic skin can be used to simulate an oesophageal or duodenal atresia repair. Manufactured anatomical models using 3D printing technology have widespread application in undergraduate teaching (Fig. 1) [11–14]. These are produced with such a high level of realism that they are often indistinguishable from biologic dissections. In a randomised controlled trial of medical students, 3D prints were found to be superior to cadaveric materials for learning external cardiac anatomy [12], perhaps because of increased physical interaction by learners. Additive manufacturing technology is advancing at a rapid rate with promising applications in understanding of complex congenital abnormalities. With the recent introduction of composite and biological printers, *operable* prints of malleable tissue become possible.

It is important to note that these PTTs are not necessarily complex. Construction should reflect fidelity as per Dieckmann [15]. The question of purpose is crucial in simulator design: what is to be learnt from the activity? Conceptual and experiential fidelity is more impactful that



Figure 1 Additive manufacturing (3D Printing) anatomical models (courtesy of Prof. P. McMnamin).




Figure 2 eoSim.

the physical fidelity. A simulator needs to confer the skills required, rather than achieve high visual reality.

Laparoscopic bench trainers

Laparoscopic surgery has been widely embraced because of the advantages of decreased post-operative pain and shorter hospital stays while offering equivalent outcomes to open surgery in many conditions. However, laparoscopic surgery involves a different skill set to traditional open surgical interventions. In particular, it requires [16]:

- Altered hand-eye co-ordination, as there is no direct vision of the operative field.
- 2D-to-3D perception realization, with the surgeon converting a 2D screen image to the 3D conceptual environment in which they operate.
- Increased fine motor skills as small hand movements are amplified by longer instruments.
- Adaptation to the fulcrum effect of the patient's body wall, causing hand movements to the left of the patient's body to generate instrument movement to the right of the operative field.
- Loss of haptic feedback as well as increased distance of surgeon from target tissues.

These vital skills are generic for all surgical specialities and can be acquired prior to patient contact. There is a developing evidence base that training on a laparoscopic bench trainer decreases the time taken to perform a laparoscopic task, improves operative accuracy, decreases errors, and improves overall performance for trainees with no previous experience in real laparoscopic surgery [17].

Bench trainers are ideally suited for application of both mastery learning and deliberate practice. They allow competency of participants to be measured against the benchmark of an expert. Through targeted feedback, improvement of technique toward competence is facilitated. Independent deliberate practice of a specific skill or activity should be coupled with supported sessions with an expert trainer surgeon to provide targeted progression

through the various steps to achieve competency. In addition to a formative role in skill acquisition and refinement, bench trainers have been used in summative assessment for certification [18]; such as in the well-established Fundamentals of Laparoscopic Surgery (FLS) [19] programme, developed by the Society of American Gastrointestinal and Endoscopic Surgeons and endorsed by the American College of Surgeons [18]. This programme includes a didactic component presented in a standardised fashion, a simulation-based technical skills component, and a cognitive and technical skills assessment component.

Traditionally, bench trainers were expensive and often only accessible in dedicated training centres. Recently, new and cheaper trainers have become available [20–23], such as the eoSim trainer (eoSurgical, Edinburgh, UK) (Fig. 2). This trainer has proven construct, concurrent, and content validity against more established bench trainers [21]. It also offers access to an online curriculum via in-built motion instrument tracking that allows home-based trainee-driven deliberate practice [24]. Reducing equipment costs allows laparoscopic bench trainers to be used in low-resource settings [18,25–27]. Costs of simulation programmes for early skills development can be offset by decreased operative time as well as reduction in operative complication rates [28–31].

Virtual reality trainers

Virtual reality trainers (VR-T) have similar purpose and underpinning principles to laparoscopic bench trainers, albeit with greater sophistication of technological interface. The vastly increased cost often limits accessibility to training involving these devices. Examples include LapSim (Surgical Science, Göteborg, Sweden) (Fig. 2), and LAP Mentor (Symbionix, Cleveland, OH, USA). Computer-based VR-Ts may be used for both basic and advanced laparoscopic skills training, with an evolving evidence base for use in a surgical curriculum [32].

VR-T use is seen most widely in the field of robotic surgery, supporting its exponential growth in recent years. There are multiple trainers available for robotic platforms, such as dV-Trainer (Mimic Technologies, Inc, Seattle, WA), the Robotic Surgical Simulator (RoSS; Simulated Surgical Systems, Buffalo, NY), and the SEP Robot (SimSurgery, Oslo, Norway) [33]. Some systems require training prior to clinical use, even for expert laparoscopic surgeons.

The safety of the VR environment for skills acquisition is both an advantage and a concern; with potential for creating complacency and false confidence that may translate to the operating theatre environment. The inherent disconnect from consequences of errors and complications can be overcome with use of hybrid simulation to add layers of complexity to the activity and task.

Simulated patients and role-play

Simulated patients (SPs) are widely used in medical education [35–37], driven by the need to enhance communication and patient-contact skills. SPs are particularly valuable in standardised assessment settings such as OSCEs (objective standardised clinical examinations). SPs are

trained to represent patients or their relatives in a simulated clinical encounter. Detailed role descriptions ensure 'patient' responses are standardised [35,38] to allow assessment of interpersonal and professional skills of a learner. In some circumstances, specific SP feedback to the learner can be very informative [35]. Often, professional SPs will work and study with real clinical patients to improve their responses and therefore improve the quality of the simulation encounter.

SPs may be involved in both formative and summative activities; promoting development of non-technical skills as well as providing a method by which to assess learner performance in these domains. In addition, SPs may be used in hybrid and scenario-based simulation activities.

Role-play is a widely used educational method for learning about communication skills in medical education. It essentially focuses on inter-personal interaction, with the participants ideally acting in their normal roles, although it can be used to change attitudes by acting in different ones. These activities often include the use of SPs as this enables standardisation of the process and, therefore, it can also be used as an assessment tool. Specific examples include obtaining consent from a parent prior to a urological intervention, informing a parent of an intra-operative complication, or breaking bad news.

Hybrid simulation

Hybrid simulation involves the pairing of PTTs with SPs or other simulation modalities. The combination aims to simulate the complexity of real-world clinical scenarios, enabling the learner to develop or be assessed on multiple skills concurrently [39]. This is often conducted *in situ*, for example emergency department resuscitation bay, outpatient department, or clinical ward to add further contextual reality. An example is a latex skin laceration simulator positioned on a SP arm (Fig. 3) [40]. As in the real-world environment, the learner needs to suture the laceration while interacting with the SP, requiring integration of technical skills with effective communication skills [41]. In urologic application, models are available to allow acquisition of urethral catheterisation technique, while simultaneously interacting with the patient. A more advanced



Figure 3 Hybrid simulation used in a distributed simulation environment [40].

example involves simulated flexible cystoscopy and stent removal, employing a fully draped PTT in a simulated operating theatre requiring communication with both nurse and SP [42]. Complexity could be added with anaesthetic complications, poor support from a nurse, a distressed or uncooperative patient, or an adherent stent which is unable to be removed. Similarly, a laparoscopic nephrectomy model, which simulates acute blood loss, has been used in a urological training capacity to assess crisis management and communication skills within a surgical team using a laparoscopic bench trainer in an operating theatre environment [43].

Scenario-based simulation

Scenario-based simulation is a widely applied form of SBME in surgical training, forming the core of many essential courses such as Advanced Trauma Life Support (ATLS) and Advanced Paediatric Life Support (APLS). Although scenarios are used for practical and technical skills acquisition in this setting, it can be used for development of non-procedural skills such as communication skills, situational awareness, expert judgement, decision-making skills, and cognitive task analysis [44]. The aim of a scenario-based simulation is to develop an instructionally sound activity that fulfils the learning objectives of the session. Scenario development requires a timeline for crucial events and variations in patient observations. Video debriefing is often used for this form of simulation, but should be conducted by adequately trained facilitators to provide safety for the learners [45–47].

The importance of feedback and discussion in learning from scenario-based simulation cannot be overemphasized. The promotion of reflective practice in experiential learning is vital. Complex emotions can be involved, especially where participants have prior clinical experience with poor outcomes.

Scenarios can be conducted within a purpose-built simulation centre (Fig. 4) or *in situ* within a clinical operating theatre, ward, or emergency department environment. Non-technical skills conferred in SBME team-based training can potentially improve patient outcome, although the evidence base for this is still being established [48]. Examples of scenario-based learning include:

cardiovascular arrest of a patient on the ward following a laparoscopic pyeloplasty, communication skills with junior theatre staff during complex hypospadias repair, or dealing with an angry and upset parent as they present with a urethral fistula post-operatively in the clinic.

Distributed simulation

Distributed simulation is a disruptive innovation that involves creation of a high-fidelity immersive environment in a portable and inexpensive format, allowing use whenever and wherever it is required [40]. Permitting a simulation team to function in any educational space increases the accessibility to learners while minimizing disruption to local clinical service provisions. Key elements of a real clinical setting are identified and recreated within a contained space, such as an inflatable structure containing a portable operating light, table, drapes, and photographic banners to recreate anaesthetic machines and trolleys (Fig. 5). A pre-recorded audio track is used to enhance experience, recreating authentic background sounds.

This environment has been combined with other simulation modalities including hybrid and scenario-based activities involving non-operative skills such as communication and team working [34]. Advantages of location




Figure 5 Operating theatre created using distributed simulation [40].



Figure 4 Operating theatre, Monash Children's Hospital Surgical Simulation Centre.

flexibility and reduced resource requirements enhance accessibility and offset disadvantages of reduced quality of video-based debriefing and feedback. Before the introduction of this disruptive innovation, high-fidelity immersive simulation was only possible in a dedicated simulation centre. Potential future applications in low-resource settings and countries exist, but have not yet been subject to trial.

Virtual reality and online simulation

Virtual reality simulation has gained popularity in recent years, with examples including interprofessional and communication skills learning through virtual human experience [49], assessment of fine motor skills prior to a clinical encounter [50], and team training in an operating theatre [51]. Links to interactive gaming environments are clear in examples such as Zero Hour: America's Medic, a first-person video game designed to train first responders to mass casualty incidents [52]. While concerns about cost and false confidence exist, the interaction provides a good introduction to the principles of triage and first-line patient management.

Online learning resources are becoming more widespread in paediatric surgical and urological training. Resources such as WeBSurg allow both trainees and practicing surgeons to view procedures prior to clinical contact [53]. This has been incorporated in the formation of a Global Virtual University [54], aiding the formation of virtual surgical communities of practice [55]. A urological example of online resources is the CEVL interactive system [56]; in this online educational module, the entire operating room staff are able to train with the system and hence improve performance with a paediatric robotic-assisted laparoscopic pyeloplasty.

Conclusions

There are many different modalities of simulation applicable to surgical training. Some of these are readily available and other are more expensive; located in specialised surgical simulation centres. Advances in simulator design have resulted in decreased cost and bring aspects of SBME within the reach of all departments, even in resource-constrained environments. Surgical simulation does not have to be expensive or time-consuming to be effective and may therefore be incorporated into busy surgical timetables. Accessibility promotes self-directed learning by the surgical trainee but this needs to be augmented with regular observation and feedback from an expert trainer.

Regardless of the technology used in educational activities, the principle focus must be on the purpose or learning intention. Simulation does not have to recreate an exact clinical environment or procedure to be useful; however, high conceptual and experiential fidelity is important. Technologically advanced mannequins and simulators may be imagined offering greater fidelity, but this is not always true. A simple hook on a board has an excellent conceptual fidelity if the purpose is for teaching hand-tying of knots. The drive for more technology to be incorporated into simulation needs to be balanced with the purpose of the

learning experience. With all simulation-based education, clarification of learning objectives for an activity can aid identification of the most appropriate modality to achieve these.

Conflict of interest

None.

Funding

None.

References

- [1] Aggarwal R, Darzi A. Technical-skills training in the 21st century. *N Engl J Med* 2006 Dec 21;355(25):2695–6.
- [2] Flin R, Yule S, Paterson-Brown S, Maran N, Rowley D, Youngson G. Teaching surgeons about non-technical skills. *Surgeon* 2007 Apr 1;5(2):86–9.
- [3] Gawande AA, Zinner MJ, Studdert DM, Brennan TA. Analysis of errors reported by surgeons at three teaching hospitals. *Surgery* 2003 Jun;133(6):614–21.
- [4] Christian CK, Gustafson ML, Roth EM, Sheridan TB, Gandhi TK, Dwyer K, et al. A prospective study of patient safety in the operating room. *Surgery* 2006 Feb;139(2):159–73.
- [5] Owen H. Early use of simulation in medical education. *Simul Healthc* 2012 Apr 1;7(2):102–16.
- [6] Fonseca AL, Evans LV, Gusberg RJ. Open surgical simulation in residency training: a review of its status and a case for its incorporation. *J Surg Educ* 2013 Jan;70(1):129–37.
- [7] Anastakis DJ, Regehr G, Reznick RK, Cusimano M, Murnaghan J, Brown M, et al. Assessment of technical skills transfer from the bench training model to the human model. *Am J Surg* 1999 Feb;177(2):167–70.
- [8] Hamdorf JM, Hall JC. Acquiring surgical skills. *Br J Surg* 2000 Jan 1;87(1):28–37.
- [9] Barsness KA, Rooney DM, Davis LM, O'Brien E. Evaluation of three sources of validity evidence for a laparoscopic duodenal atresia repair simulator. *J Laparoendosc Adv Surg Tech A* 2015 Mar;25(3):256–60.
- [10] Barsness KA, Rooney DM, Davis LM, O'Brien E. Evaluation of three sources of validity evidence for a synthetic thoracoscopic esophageal atresia/tracheoesophageal fistula repair simulator. *J Laparoendosc Adv Surg Tech A* 2015 Jul;25(7):599–604.
- [11] McMenamin PG, Quayle MR, McHenry CR, Adams JW. The production of anatomical teaching resources using three-dimensional (3D) printing technology. *Anat Sci Educ* 2014 Nov 12;7(6):479–86.
- [12] Lim KHA, Loo ZY, Goldie SJ, Adams JW, McMenamin PG. Use of 3D printed models in medical education: a randomized control trial comparing 3D prints versus cadaveric materials for learning external cardiac anatomy. *Anat Sci Educ* 2016 May 6;9(3):213–21.
- [13] Lioufas PA, Quayle MR, Leong JC, McMenamin PG. 3D printed models of cleft palate pathology for surgical education. *Plast Reconstr Surg Glob Open* 2016 Sep;4(9):e1029.
- [14] Loke Y-H, Harahsheh AS, Krieger A, Olivieri LJ. Usage of 3D models of tetralogy of Fallot for medical education: impact on learning congenital heart disease. *BMC Med Educ* 2017 Mar 11;17(1):54.
- [15] Dieckmann P, Gaba D, Rall M. Deepening the theoretical foundations of patient simulation as social practice. *Simul Healthc* 2007;2(3):183–93.

- [16] Gallagher AG, McClure N, McGuigan J, Crothers I, Browning J. Virtual reality training in laparoscopic surgery: a preliminary assessment of minimally invasive surgical trainer virtual reality (MIST VR). *Endoscopy* 1999 May;31(4):310–3.
- [17] Nagendran M, Toon CD, Davidson BR, Gurusamy KS. Laparoscopic surgical box model training for surgical trainees with no prior laparoscopic experience. *Cochrane Database Syst Rev* 2014 Jan 17;(1), CD010479.
- [18] Okrainec A, Soper NJ, Swanstrom LL, Fried GM. Trends and results of the first 5 years of Fundamentals of Laparoscopic Surgery (FLS) certification testing. *Surg Endosc* 2011;25(4):1192–8.
- [19] Peters JH, Fried GM, Swanstrom LL, Soper NJ, Sillin LF, Schirmer B, et al. Development and validation of a comprehensive program of education and assessment of the basic fundamentals of laparoscopic surgery. *Surgery* 2004 Jan;135(1):21–7.
- [20] Van Sickle KR, McClusky III DA, Gallagher AG, Smith CD. Construct validation of the ProMIS simulator using a novel laparoscopic suturing task. *Surg Endosc* 2005;19(9):1227–31.
- [21] Hennessey IAM, Hewett P. Construct, concurrent, and content validity of the eoSim laparoscopic simulator. *J Laparoendosc Adv Surg Tech A* 2013 Oct;23(10):855–60.
- [22] Martinez AM, Espinoza DL. Novel laparoscopic home trainer. *Surg Laparosc Endosc Percutan Tech* 2007 Aug 1;17(4):300.
- [23] Rivas AM, Vilanova AC, Pereferrer FTS, Gonz lez MHN, del Castillo Déjardin D. Low cost simulator for acquiring basic laparoscopic skills. *Cir Esp (English Edition)* 2010 Jan;87(1):26–32.
- [24] Partridge RW, Hughes MA, Brennan PM, Hennessey IAM. Accessible laparoscopic instrument tracking (“InsTrac”): construct validity in a take-home box simulator. *J Laparoendosc Adv Surg Tech A* 2014 Aug;24(8):578–83.
- [25] Kolozsvari NO, Kaneva P, Vassiliou MC, Fried GM, Feldman LS. New dog, new tricks: trends in performance on the Fundamentals of Laparoscopic Surgery simulator for incoming surgery residents. *Surg Endosc* 2012 Jan;26(1):68–71.
- [26] Okrainec A, Smith L, Azzie G. Surgical simulation in Africa: the feasibility and impact of a 3-day fundamentals of laparoscopic surgery course. *Surg Endosc* 2009 Nov;23(11):2493–8.
- [27] Rosenthal ME, Ritter EM, Goova MT, Castellvi AO, Tesfay ST, Pimentel EA, et al. Proficiency-based Fundamentals of Laparoscopic Surgery skills training results in durable performance improvement and a uniform certification pass rate. *Surg Endosc* 2010 Oct 1;24(10):2453–7.
- [28] Farnworth LR, Lemay DE, Wooldridge T, Mabrey JD, Blaschak MJ, DeCoster TA, et al. A comparison of operative times in arthroscopic ACL reconstruction between orthopaedic faculty and residents: the financial impact of orthopaedic surgical training in the operating room. *Iowa Orthop J* 2001;21:31–5.
- [29] Wilkiemeyer M, Pappas TN, Giobbie-Hurder A, Itani KMF, Jonasson O, Neumayer LA. Does resident post graduate year influence the outcomes of inguinal hernia repair? *Ann Surg* 2005 Jun 1;241(6):879–84.
- [30] Kauvar DS, Braswell A, Brown BD, Harnisch M. Influence of resident and attending surgeon seniority on operative performance in laparoscopic cholecystectomy. *J Surg Res* 2006 May;132(2):159–63.
- [31] Bridges M, Diamond DL. The financial impact of teaching surgical residents in the operating room. *Am J Surg* 1999 Jan;177(1):28–32.
- [32] Beyer-Berjot L, Berdah S, Hashimoto DA, Darzi A, Aggarwal R. A virtual reality training curriculum for laparoscopic colorectal surgery. *J Surg Educ* 2016 Dec;73(6):932–41.
- [33] Lallas CD, Davis JW, Members Of The Society Of Urologic Robotic Surgeons. Robotic surgery training with commercially available simulation systems in 2011: a current review and practice pattern survey from the society of urologic robotic surgeons. *J Endourol* 2012 Mar;26(3):283–93.
- [34] Kassab E, Tun JK, Arora S, King D, Ahmed K, Miskovic D, et al. “Blowing up the barriers” in surgical training: exploring and validating the concept of distributed simulation. *Ann Surg* 2011 Dec 1;254(6):1059–65.
- [35] Cleland JA, Abe K, Rethans J-J. The use of simulated patients in medical education: AMEE Guide No 42. *Med Teach* 2009 Aug 27;31(6):477–86.
- [36] Hargie O, Dickson D, Boohan M, Hughes K. A survey of communication skills training in UK Schools of Medicine: present practices and prospective proposals. *Med Educ* 1998 Jan 1;32(1):25–34.
- [37] Nestel D. *Simulated patient methodology*. Chichester, UK: John Wiley & Son’s; 2014 Nov 3. p. 1–9.
- [38] Ker JS, Dowie A, Dowell J, Dewar G, Dent JA, Ramsay J, et al. Twelve tips for developing and maintaining a simulated patient bank. *Med Teach* 2009 Jul 3;27(1):4–9.
- [39] Kjellin A, Hedman L, Escher C, Felländer-Tsai L. Hybrid simulation: bringing motivation to the art of teamwork training in the operating room. *Scand J Surg* 2014 Dec;103(4):232–6.
- [40] Kneebone R, Arora S, King D, Bello F, Sevdalis N, Kassab E, et al. Distributed simulation – accessible immersive training. *Med Teach* 2010 Jan 22;32(1):65–70.
- [41] Kneebone R, Kidd J, Nestel D, Asvall S, Paraskeva P, Darzi A. An innovative model for teaching and learning clinical procedures. *Med Educ* 2002 Jul 1;36(7):628–34.
- [42] Nguyen LN, Tardioli K, Roberts M, Watterson J. Development and incorporation of hybrid simulation OSCE into in-training examinations to assess multiple CanMEDS competencies in urologic trainees. *Can Urol Assoc J* 2015 Jan;9(1–2):32–6.
- [43] Lee JY, Mucksavage P, Canales C, McDougall EM, Lin S. High fidelity simulation based team training in urology: a preliminary interdisciplinary study of technical and nontechnical skills in laparoscopic complications management. *J Urol* 2012 Apr;187(4):1385–91.
- [44] Andersen DK. How can educators use simulation applications to teach and assess surgical judgment? *Acad Med* 2012 Jul 1;87(7):934.
- [45] Cheng A, Eppich W, Grant V, Sherbino J, Zendejas B, Cook DA. Debriefing for technology-enhanced simulation: a systematic review and meta-analysis. *Med Educ* 2014 Jun 9;48(7):657–66.
- [46] Krogh K, Bearman M, Nestel D. “Thinking on your feet”—a qualitative study of debriefing practice. *Adv Simul* 2016 Jun 15:1–11.
- [47] Rudolph JW, Raemer DB, Simon R. Establishing a safe container for learning in simulation. *Simul Healthc* 2014 Dec;9(6):339–49.
- [48] Owei L, Neylan CJ, Rao R, Caskey RC, Morris JB, Sensenig R, et al. In situ operating room? Based simulation: a review. *J Surg Educ* 2017 Mar;0(0).
- [49] Johnsen K, Raij A, Stevens A, Lind DS. The validity of a virtual human experience for interpersonal skills education. In: CHI’07 Proceedings of the SIGCHI conference on human factors in computing systems; 2007. p. 1049–58. April 28–May 03 2007.
- [50] Mohamadipanah H, Parthiban C, Nathwani J, Rutherford D, DiMarco S, Pugh C. Can a virtual reality assessment of fine motor skill predict successful central line insertion? *Am J Surg* 2016 Oct;212(4):573–578.e1.
- [51] Abelson JS, Silverman E, Banfelder J, Naides A, Costa R, Dakin G. Virtual operating room for team training in surgery. *Am J Surg* 2015 Sep;210(3):585–90.
- [52] Zero Hour: America’s Medic [Internet]. Wikipedia. Available from: <http://en.wikipedia.org/wiki/Zero%20Hour%3A%20America%27s%20Medic>.
- [53] Mutter D, Vix M, Dallemagne B, Perretta S, Leroy J, Marescaux J. WebSurg: an innovative educational Web site in

- minimally invasive surgery—principles and results. *Surg Innov* 2011 Mar;18(1):8–14.
- [54] Mutter D, Rubino F, Temporal MSG, Marescaux J. Surgical education and internet-based simulation: the World Virtual University. *Minim Invasive Ther Allied Technol* 2005;14(4): 267–74.
- [55] Gandamihardja TAK. The role of communities of practice in surgical education. *J Surg Educ* 2014 Jul;71(4):645–9.
- [56] Li B, Maizels M, Liu D, Gong EM-L, Lindgren BW, Malhotra N, et al. CEVL interactive – promoting effective teamwork to perform robot assisted laparoscopic pyeloplasty in pediatric urology. *J Pediatr Urol* 2016 Dec;12(6):430–3.